

University of Dundee

Risk factors for neuropathic pain in diabetes mellitus

Hébert, Harry L.; Veluchamy, Abirami; Torrance, Nicola; Smith, Blair H.

Published in:
Pain

DOI:
[10.1097/j.pain.0000000000000785](https://doi.org/10.1097/j.pain.0000000000000785)

Publication date:
2017

Licence:
CC BY-NC-ND

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Hébert, H. L., Veluchamy, A., Torrance, N., & Smith, B. H. (2017). Risk factors for neuropathic pain in diabetes mellitus. *Pain*, 158(4), 560-568. <https://doi.org/10.1097/j.pain.0000000000000785>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Risk factors for neuropathic pain in diabetes mellitus

Harry L. Hébert*, Abirami Veluchamy, Nicola Torrance, Blair H. Smith

1. Introduction

According to the International Diabetes Federation, diabetes mellitus (DM) is estimated to affect around 415 million adults worldwide, roughly 8.8% of the adult population, with the figure projected to rise to over 600 million by 2040.²⁰ Regional prevalence varies from 3.2% in Africa to 12.9% in North America. Diabetes mellitus is associated with a number of chronic sequelae and around 50% of people with DM go on to develop polyneuropathy.³⁵ This condition has a variety of clinical manifestations, which are grouped into positive symptoms including dysesthesia (abnormal sense of touch), tingling and itching, and negative symptoms including numbness, muscle weakness, and trouble with balance. Up to 25% of people with diabetic neuropathy (DN) also develop neuropathic pain (NP).³⁹ Neuropathic pain is defined by the International Association for The Study of Pain as “pain directly caused by a lesion or disease affecting the somatosensory system.”^{12,22} Symptoms of painful diabetic neuropathy (PDN) include those described above for nonpainful DN with additional “burning,” “electric shocks,” “stabbing,” and “pins and needles” symptoms all being described. Painful diabetic neuropathy is associated with increased distress and poor quality of life compared with nonpainful DN, DM, and the general population³⁸ including depression, anxiety, and sleep disturbance.¹⁵ In addition, an association has been described with reduced productivity and employability at work compared with nonpainful DN.³⁷ The combination of these factors places a large economic burden on patients and health care services,¹⁰ a situation likely to grow steadily worse with the aforementioned projected rise in DM prevalence. This situation is further exacerbated by the fact that 13% of patients with PDN do not report their symptoms to primary care, and 39% of patients with PDN have never received treatment.⁸ Even for those patients who do attend primary and secondary care for their diabetes, pain

is not a symptom that is always included in clinical assessments. Furthermore, not all patients with DN develop PDN, and the reasons for this are unclear.

Understanding the risk factors for PDN will go some way to resolving this and will also help to inform management and prevention of this painful condition by health care services. Any factor that increases the risk of DM or DN is likely to be a risk factor for PDN. However, it is the specific nature and magnitude of the risk that remains unclear and is the focus of this topical review.

2. Risk factors

There have been relatively few published studies examining risk factors specifically for PDN in DM. Clinical, environmental, and genetic factors have been shown to be predictive of developing DM and some of these have also been implicated in the development of DN, including age,¹¹ body mass index,^{25,28} hypertension,¹³ smoking, and waist circumference²⁸ (Fig. 1). Given the likely overlap of risk factors between DM and DN, it seems reasonable to hypothesize that some of these factors will also influence the development of PDN.

We conducted a literature search using relevant key words and terms aiming to identify a wide range of studies that investigated risk factors for PDN and to include all the important studies (Table 1). A number of limitations can be identified with these studies as a whole. Most of these studies are cross sectional in nature and therefore unable to establish temporal relationship between patient characteristics/factors and PDN. Some studies report only univariate analysis and are therefore unable to assess intervariable relationships and to identify confounding between variables.^{2,6,8,9,17,23,30} In addition, it is not always clear in the methods and statistical analyses whether PDN or nonpainful DN is being analysed and what control group the PDN subjects are being compared with. In some studies, those in the control group are diabetic participants with nonpainful neuropathy^{30,38,41} and in others they are diabetic participants without neuropathy of any form.^{1,2,6,8,11,17} In other studies, it is not possible to determine the nature of the control group from the description of the methods. There was considerable heterogeneity in PDN case ascertainment, with only 6 studies using a validated NP screening questionnaire (the DN4^{6,7,17,21,38} or the Leeds assessment of neuropathic symptoms and signs¹¹) with the remainder using nonvalidated questionnaires or clinical examinations. This makes it difficult to assess the sensitivity and specificity of each study to identify PDN cases and to make direct comparisons between studies as effect size estimations and associations are likely to be different. Despite these limitations, some potential risk factors have emerged, including environment, clinical, lifestyle, and genetic factors.

Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

Chronic Pain Research Group, Division of Population Health Sciences, Ninewells Hospital and Medical School, Mackenzie Building, University of Dundee, Dundee, Scotland, United Kingdom

*Corresponding author. Address: Chronic Pain Research Group, Division of Population Health Sciences, Ninewells Hospital and Medical School, Mackenzie Building, University of Dundee, Dundee DD2 4BF, Scotland, United Kingdom. Tel.: + (44) 01382 383191. E-mail address: h.hebert@dundee.ac.uk (H. L. Hébert).

PAIN 158 (2017) 560–568

Copyright © 2016 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the International Association for the Study of Pain. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

<http://dx.doi.org/10.1097/j.pain.0000000000000785>




Figure 1. Schematic of the process from diabetes mellitus to diabetic neuropathy and finally painful diabetic neuropathy. Both diabetes mellitus and diabetic neuropathy have their own set of risk factors, both of which could provide important clues as to the risk factors that contribute to painful diabetic neuropathy.

2.1. Demographic

Two nonmodifiable factors—age^{2,17,21,38,42} and sex^{1,11,17,21,41}—have been specifically associated with PDN, in addition to their known roles as risk factors for DM. Although these are of limited use to clinicians in terms of intervention, they could provide useful clues as to the underlying biological pathways involved and increased awareness of at-risk patients. In particular, the association of PDN with older age (>50 years) is likely to be related to the time it takes for nerve damage and painful symptoms to develop after the onset of DM and the decreased ability of the body to deal with this. Similarly, gender associations may indicate possible subtle differences in biology and psychosocial factors that affect the risk of PDN, something that requires further investigation. It is interesting to note that while 4 of the studies report greater risk in women,^{1,11,17,21} 1 study reports greater risk in men.⁴¹ This discrepancy in the latter study could be related to the limited statistical analysis, which did not adjust for potential confounding factors. Despite the prevalence of DM varying according to ethnicity, this has not been found as a risk factor so far for PDN.^{1,17,18,21,32,36} One study reported that South Asians were more likely to report painful symptoms than people in other ethnic groups in the absence of clinical neuropathy.¹ Another found an association with pain among people with DM residing in a Gulf state and Lebanon compared with Egypt, but did not analyse ethnic origin.²¹

2.2. Clinical

Clinical and physiological factors associated with PDN are important for clinicians and primary care as they may indicate possibilities for targeted treatment or primary prevention strategies. The clinical diagnosis of the type of DM and the duration since onset of the disease may be particularly relevant. Two studies found an association with DM type in multivariate analysis, with 1 identifying type 1 diabetes (T1D)²¹ and the other type 2 diabetes (T2D)¹ as conferring greater risk of PDN. Differences in case definition and study populations could have contributed to the heterogeneity in these results. A clearer consensus is apparent for DM duration with risk increasing over time since diagnosis.^{2,6,17,21,32,38} Severity of preceding neuropathy has been found

to be associated with PDN, but associations with neuropathy duration and comparison with type of (peripheral or sensory) neuropathy have not been found.^{9,11,33,36} Most studies included only 1 type of neuropathy in their analysis. A number of clinical factors and comorbidities have been found to be associated with PDN. These include poor glycaemic control and high HbA1c levels,^{2,18,36} hypertension,^{2,18} retinopathy,² nephropathy,^{2,38} cardiovascular disease,^{2,42,43} and glycosuria.¹⁸ However, as these conditions are all known complications of DM, it is uncertain from cross-sectional analysis whether these factors are contributing to PDN risk and onset, or simply coexisting factors, perhaps confounded by other factors or with shared aetiology. Biomarkers for the development of PDN can be exploited by providing preventative or diagnostic tests. In this respect, tumour necrosis factor alpha and inducible nitric oxide synthase expression,³⁰ triglycerides, and low high-density lipoprotein cholesterol³⁸ all show promising associations but require replication to be confident in their role in disease pathogenesis.

2.3. Lifestyle

Behavioural and social circumstances are important lifestyle aspects that patients can theoretically influence and act on, with greater or less practical difficulty. In particular, some physical characteristics known to be associated with DM and DN are also implicated in PDN. Body mass index has been clearly linked to PDN, particularly in the form of obesity ($\geq 30 \text{ kg/m}^2$),^{21,33,38} while in another study, weight was reported independently of height and found to be significantly associated with PDN, although this was attenuated in multivariate analysis.⁴² A related study also found a positive correlation with increased waist circumference and high levels of physical activity and risk of PDN.⁴³ Despite being included in the analyses in most of the studies,^{1,2,4,8,17,18,21,32,33,38,42,43} smoking and alcohol consumption have not been specifically associated with PDN. Psychological factors have also been widely reported in the context of general chronic pain, but its relationship with PDN is less clear. Increased depression, anxiety, enjoyment of life, and social relationships are associated with PDN, but without prospective studies and longitudinal analysis, the temporal relationship cannot be established.^{7,15}

Table 1

List of studies conducted on predictors of painful diabetic neuropathy and their characteristics.

Reference	Study type	Population	Criteria for NP	Sample size	Analysis	Variables analysed	Predictors	OR (95% CI)/P
Abbott et al., 2011 ¹	Cross sectional	UK	NSS ≥ 5 and NDS ≥ 3	3242 DM with NP	Multivariate logistic regression	Age, alcohol, diabetes duration, diabetes treatment, diabetes type, ethnicity, foot ulcer, foot deformities, impaired vision, lower limb amputation, nephropathy, PAD, sex, and smoking	T2D	2.1 (1.7-2.4)
				12,372 DM without PDN			Women	1.5 (1.4-1.6)
AlQuliti, 2015 ²	Case control	Saudi Arabia	Foot examination and NSS ≥3	99 T2D with PDN	Univariate analysis	Age, CVD, diabetes duration, glycaemic control, HbA1c, hypertension, insulin use, nephropathy, oral antidiabetic drugs, PVD, retinopathy, sex, smoking, stroke, and working status	Age (>50 y)	1.93 (1.09-3.41)
				99 T2D without PDN			CVD	3.37 (1.28-8.89)
							Diabetes duration (>10 y)	3.38 (1.88-6.07)
							Glycaemic control	0.42 (0.12-0.96)
							Hypertension	2.85 (1.57-5.17)
							Nephropathy	8.93 (2.58-30.95)
							Retinopathy	13.22 (4.49-38.95)
Benbow et al., 1997 ⁴	Cross sectional	UK	Clinical history and examination; burning/shooting pain/hyperesthesia ≥6 mo and at least 1 abnormal neurological sign from decrease in light touch, vibration, or pinprick sensation	49 DN with NP	Univariate analysis	Age, diabetes duration, diabetes type, HbA1c, sex, and smoking	NA	NA
				23 DN without NP				
Cheng et al., 2010 ⁵	Genetic case-control/cross sectional	Taiwan	Pain VAS ≥4 and grade 3-5 of occurrence of pain in daily activities	15 DFU (and DN) with NP	Univariate analysis/Fisher exact	Age, albumin, amputation, BMI, diabetes duration, diabetes type, haemoglobin, HbA1c, hyperlipidemia, hypertension, rs1799971 of <i>OPRM1</i> , and sex	rs1799971	0.24 (0.07-0.8)
				50 DFU (and DN) without NP				
Cortez et al., 2014 ⁶	Cross sectional	Brazil	DN4 ≥ 4	12 T2D with PDN	Multivariate analysis	Age, depressive symptoms, diabetes duration, drug adhesion, sex, and glycaemic control	Diabetes duration	P = 0.031
				60 T2D without PDN				
D'Amato et al., 2016 ⁷	Cross sectional	Italy	DN4 ≥ 4 (DN4 interview ≥ 3)	25 DN with NP 46 DN without NP 110 without DN	Multivariate analysis	Depression	Depression (BDI-II)	4.56 (1.09-19.1)

(continued on next page)

Table 1 (continued)

Reference	Study type	Population	Criteria for NP	Sample size	Analysis	Variables analysed	Predictors	OR (95% CI)/P
Daousi et al., 2004 ⁸	Cross sectional	UK	Typical NP symptoms in legs ≥ 1 y, PSS ≥ 3 and NDS ≥ 6 or NDS ≥ 3 and NSS ≥ 5	56 DM with PDN	Univariate analysis	Age, alcohol, angina, BMI, BP, CVA, depression, diabetes duration, diabetes type, foot ulceration, HbA1c, hypertension, MI, PVD, sex, and smoking	NA	NA
				289 DM without PDN				
Davies et al., 2006 ⁹	Cross sectional	UK	Positive response to "Do you have a burning, aching or tenderness in your legs or feet?" from DNSS and TCSS score >5	71 T2D with PDN (51 with NP and 20 with mixed NP and non-NP) 99 T2D with non-NP 99 T2D with no pain	Univariate analysis	Age, diabetes duration, HbA1c, neuropathy severity, and sex	Severity of neuropathy	$P < 0.0001$
Erbas et al., 2011 ¹¹	Cross sectional	Turkey	LANSS ≥ 12	156 DM with PDN	Univariate analysis	Age, blood urea BMI, BUN, creatinine, diabetes duration, diabetes type, FPG, HbA1c, PPG, and sex	Duration of diabetes	$P = 0.001$
				975 DM without PDN			T1D Women	$P = 0.039$ $P = 0.001$
Gore et al., 2005 ¹⁵	Cross sectional	USA	Physician diagnosed	255 with PDN	Univariate analysis	Anxiety, depression, enjoyment of life, mental health, mood, and relationship with others	Anxiety (HADS) Depression (HADS) Enjoyment of life (BPI-DPN) Mental health (SF-12v2) Mood (BPI-DPN) Relationship with others (BPI-DPN)	All $P < 0.05$
Halawa et al., 2010 ¹⁷	Cross sectional	Saudi Arabia	DN4 ≥ 4	678 DM with PDN	Univariate analysis	Age, BMI, diabetes duration, diabetes type, ethnicity, smoking, and sex	Age	$P < 0.001$
				361 DM without PDN			Diabetes duration Women	$P < 0.001$ $P = 0.024$
Harris et al., 1993 ¹⁸	Cross sectional	USA	Positive response to, "During the past 3 mo, have you had a painful sensation or tingling in your hands or feet?"	2392 with DM (26.8% of whom had pain/tingling in hands/feet?)	Multivariate logistic regression	Age, amputation, angina, diabetes age, diabetes duration, ethnicity, family income, foot sores, height, higher education, hypertension, insulin, nephropathy, obesity, periodontal disease, proteinuria, retinopathy, sex, smoking, and stroke	Glycosuria	2.31 (1.54-3.47)
				20,037 without DM			Hyperglycaemia Hypertension	2.51 (1.81-3.49) 1.58 (1.31-1.90)

(continued on next page)

Table 1 (continued)

Reference	Study type	Population	Criteria for NP	Sample size	Analysis	Variables analysed	Predictors	OR (95% CI)/P
Jambart et al., 2011 ²¹	Cross sectional	Middle East Region	DN4 \geq 4	2144 DM with PDN	Multivariate regression	Age, BMI, diabetes duration, diabetes type, ethnicity, sex, and smoking	Age (50–64 y)	1.75 (1.48–2.08)
				1845 DM without PDN			Age (\geq 65 y)	2.13 (1.72–2.62)
							BMI (\geq 30 kg/m ²)	1.35 (1.17–1.56)
							Diabetes duration (\geq 10 y)	2.43 (2.10–2.81)
							Living in a Gulf State (compared with Egypt)	0.44 (0.35–0.56)
							Living in Lebanon (compared with Egypt)	0.66 (0.54–0.81)
							T1D	1.59 (1.24–2.05)
							Women	1.27 (1.11–1.46)
Li et al., 2015 ²³	Genetic case control	USA/Canada	NCT00501202: lower extremity pain \geq 3 mo	887 DM with PDN	Univariate analysis	SNPs across <i>SCN9A</i> gene region	rs74449889 (<i>SCN9A</i>)	2.6
			NCT00870454: as above and NRS-11 \geq 11	1029 without DM and PDN			rs3750904 (<i>SCN9A</i>)	2.2
			NCT00993018: Symmetrical pain beginning in feet >6 mo and NRS-11 \geq 5 but <10 over 7 d.				rs4369876 (<i>SCN9A</i>)	2.1
			NCT00455520: clinical diagnosis with signs and symptoms >6 mo and at screening				rs12478318 (<i>SCN9A</i>)	2.1
Meng et al., 2015 ²⁷	GWAS	UK	Prescription of at least one from duloxetine, gabapentin, pregabalin, capsaicin cream/patch, and lidocaine patch. And positive monofilament test in at least 1 foot	572 DM with PDN	Fisher exact	SNPs across whole genome	rs17428041 (Chr8p21.3)	0.67 (0.57–0.78)
				2491 DM without PDN				
Meng et al., 2015 ²⁶	GWAS	UK	Multiple usage of at least 1 from duloxetine, gabapentin, pregabalin, capsaicin cream/patch, lidocaine patch	961 DM with PDN	Logistic regression	SNPs across whole genome	rs71647933 (Chr1p35.1)	2.31 (1.68–3.17)
				3260 DM without PDN			rs6986153 (Chr8p23.1)	1.67 (1.34–2.08)

(continued on next page)

Table 1 (continued)

Reference	Study type	Population	Criteria for NP	Sample size	Analysis	Variables analysed	Predictors	OR (95% CI)/P
Purwata, 2011 ³⁰	Case control	Indonesia	Pain intensity VAS >0 (representing no pain)	59 DN with NP	Univariate analysis	Age, diabetes duration, FG, HbA1c, iNOS expression, plasma TNF- α , TNF- α expression, and 2h-G	iNOS expression	3.546 (1.613-7.795)
				51 DN without NP			Plasma TNF- α	5.053 (2.241-11.392)
							TNF- α expression	4.125 (1.805-9.425)
Sorensen et al., 2002 ³²	Cross sectional	Australia	Bilateral and symmetrical foot pain—patient specifically asked about foot pain	2610 T2D (3.3% with PDN)	Multivariate logistic regression	Age, alcohol, diabetes duration, diabetes treatment, ethnicity, HbA1c, height, sex, and smoking	Diabetes Duration	1.09 (1.06-1.1)
Spallone et al., 2011 ³³	Cross sectional	Italy	Clinical examination and history	78 DN with NP	Multivariate logistic regression	Age, alcohol, BMI, BP, creatinine, CVD, diabetes duration, diabetes type, HbA1c, HDL, hypertension, LDL, nephropathy, PAD, physical activity, retinopathy, sex, smoking, triglyceride, and waist circumference	BMI (kg/m ²)	1.22 (1.08-1.37)
				57 DN without NP 56 without DN or NP			Severity of neuropathy	1.27 (1.11-1.44)
Themistocleous et al., 2016 ³⁶	Cross sectional	UK	IASP/NeuPSIG grading system	70 DPN with moderate/severe NP	Univariate analysis	Age, BMI, diabetes duration, diabetes type, ethnicity, HbA1c, neuropathy severity, orthostatic hypotension, ratio, sex, standing and lying BP, and waist–hip circumference	HbA1c and neuropathy severity	$P < 0.01$
				41 DPN with mild NP 80 DPN without NP				$P < 0.01$
Van Acker et al., 2009 ³⁸	Cross sectional	Belgium	DN4 ≥ 4 and positive Neuropen test	157 DN with NP	Multivariate logistic regression	Age, BMI, BP, diabetes duration, foot lesions, HbA1c, HDL, insulin, LDL, nephropathy, retinopathy, sex, triglycerides, and waist circumference	Age (per 10 y)	1.47 (1.20-1.81)
				321 DN without NP			Diabetes duration (per 5 y)	1.14 (1.02-1.28)
							HDL cholesterol (≤ 1 mmol/L for men, ≤ 1.3 mmol/L for women)	2.17 (1.38-3.41)
							Nephropathy	1.69 (1.10-2.59)
							Obesity (≥ 30 kg/m ²)	1.62 (1.05-2.49)
Triglycerides (≥ 1.7 mmol/L)	1.76 (1.13-2.75)							
Wu et al., 2007 ⁴¹	Cross sectional	France	MNSI ≥ 7 and Q5 of BPI ≥ 1	72 DN with NP	No statistical analysis	Age, diabetes age, diabetes duration, diabetes type, education, employment, region, and sex	Age (over 65 y)	NA
				12 DN without NP			Men	

(continued on next page)

Table 1 (continued)

Reference	Study type	Population	Criteria for NP	Sample size	Analysis	Variables analysed	Predictors	OR (95% CI)/P
Ziegler et al., 2009 ⁴²	Cross sectional	Germany	MNSI > 2 and positive answer to Q2 and Q6	195 DM (13.3% with NP)	Multivariate logistic regression	Age, albuminuria, BMI, creatinine, Fg, HbA1c, HDL, height, LDL, PAD, physical activity, sex, smoking, alcohol, stroke, systolic BP, waist circumference, and 2h-G	Age	1.08 (1.00–1.16)
Ziegler et al., 2009 ⁴³	Cross sectional	Germany	MNSI > 2 and positive answer to Q2 and/or Q6	214 DM (21.0% with NP)	Multivariate logistic regression	Age, albuminuria, BMI, creatinine, Fg, HbA1c, HDL, height, LDL, PAD, physical activity, sex, smoking, alcohol, stroke, systolic BP, waist circumference, and 2h-G	PAD Weight*	9.27 (3.44–25.0) 1.03 (1.00–1.06)*
							Physical activity Waist circumference	0.31 (0.10–0.99) 1.05 (1.01–1.09)

* Reported association, but $P > 0.05$.

2h-G, 2 hours glucose; BDI-II, Beck Depression Inventory II; BMI, body mass index; BP, blood pressure; BPI, brief pain inventory; BUN, blood urea nitrogen; CI, confidence interval; CVD, cerebrovascular disease; DM, diabetes mellitus; DN, diabetic neuropathy; DM4, Douleur Neuropathique en 4 Questions; DNSS, Diabetic Neuropathy Symptom Score; DPN, diabetic peripheral neuropathy; Fg, fasting glucose; FPG, fasting plasma glucose; GWAS, genome-wide association study; HADS, Hospital Anxiety and Depression Scale; HDL, high-density lipoprotein; IASP, International Association for the Study of Pain; INOS, inducible nitric oxide synthase; LANSS, Leeds assessment of neuropathic symptoms and signs; LDL, low-density lipoprotein; MI, myocardial infarction; MNSI, Michigan Neuropathy Screening Instrument; NA, not applicable; NDS, Neuropathy Disability Score; NeuPSIG, neuropathic pain special interest group; NP, neuropathic pain; NRS-11, numeric rating scale-11; NSS, Neuropathy Symptom Score; OR, odds ratio; PAD, peripheral arterial disease; PDN, painful diabetic neuropathy; PPG, postprandial plasma glucose; PVD, peripheral vascular disease; SCNGA, sodium voltage-gated channel alpha subunit 9; SF-12v2, Short-Form 12 version 2; SNP, single-nucleotide polymorphism; T1D, type 1 diabetes; T2D, type 2 diabetes; TCSS, Toronto clinical scoring system; TNF- α , tumour necrosis factor alpha; UK, United Kingdom; USA, United States of America; VAS, visual analogue scale.

2.4. Genetics

Numerous published studies have found that both T1D and T2D have a heritable component^{3,19,29,31} and genetic studies have been conducted in DN,^{24,34} although heritability studies have yet to be conducted. A heritable component to PDN has been hinted at in 1 study, which found that 56% of participants also reported a family member with the condition.¹⁴ Two PDN genome-wide association studies have been conducted, both in the same Scottish diabetic cohort, but with slightly differing phenotype definitions. The first used a positive monofilament test combined with a prescription history of at least 1 from 5 recommended NP medicines. This yielded an association at chromosome 8p21.3 near the gene *GFR2* and estimated the narrow-sense heritability (proportion of phenotypic variance explained by additive genetic variance) to be 11%.²⁷ The second used the same definition but without the monofilament test and aimed to capture associations that may have been missed in the previous study, using a less specific cohort. This found sex-specific associations at chromosome 1p35.1 in the *ZSCAN20/TLR12P* gene regions in females and chromosome 8p23.1 next to *HMGB1P46* in males.²⁶ The narrow-sense heritability was 30% in males and 14.7% in females. In both of these studies, controls were defined as patients who had not previously been prescribed any of the 5 NP medicines or other medicines which are predominantly prescribed for other conditions but are also known to be prescribed for NP. Two separate candidate-gene studies have been conducted and have reported associations with PDN. The first was in the sodium channel gene *SCN9A*, which is expressed in dorsal root ganglia, using a combination of numerical rating scales and clinical examinations to compare PDN with healthy controls.²³ The second was in the opioid receptor gene *OPRM1* using a visual analogue scale for pain intensity and a grading system for pain occurrence during daily activities.⁵ All 4 of these studies require independent replication, and further studies are required to establish the extent to which genetics contribute specific risk of PDN and the mechanism of this contribution.

3. Conclusions

Despite the limited number of studies reporting specific predictors for PDN, clear similarities are emerging with the known general risk factors for both DM and DN. These include clinical factors such as diabetes type and duration and lifestyle factors such as body mass index and waist circumference, some of which are not easily or not at all modifiable. Although further work is needed, this suggests that PDN is a manifestation of longer-lasting and more severe diabetes and certainly that it requires specific testing and diagnosis in routine diabetic care. However, there are likely to be factors, among those with DM (with and without DN) that confer a greater risk of PDN, and these require further exploration. It should be noted that while the published literature (and this topical review) has mainly explored NP arising from DN, NP can also arise in the diabetic population through other causes, for example, sciatic neuralgia and carpal tunnel syndrome. The influence of diabetes on the development of NP in these conditions is an area that requires further investigation. There is clear evidence to suggest PDN has a negative impact on quality of life; however, the extent to which the reverse is true—bidirectional aetiology—is currently unknown. Future studies need to be conducted in a longitudinal manner or as clinical trials to establish the temporal relationship between variable and disease, particularly with respect to identifying specific PDN risk factors in T1D and T2D patients. The previous point can be further strengthened by

running Mendelian randomization studies, something that has been used in DM.¹⁶ Mendelian randomization studies establish causal relationship by comparing 2 groups of individuals with and without a genetic marker known to influence the variable being studied. As genotype assignment is random and not subject to confounding typically found in epidemiological studies, a higher prevalence of disease in the group with the marker implies causality. However, we would first need clearer evidence to identify genetic factors associated with PDN. Finally, greater clarity is needed in specifying whether painful or nonpainful DN is being analysed. This can be enhanced by forming a consensus on PDN phenotype definition, to enable studies to be more comparable with one another. This is something that has been addressed for NP generally and could also be applied to PDN.⁴⁰ This would make replication of results more likely and brings the added potential of being able to perform meta-analyses in the future. All these limitations will be addressed in the DOLORisk study (<http://dolorisk.eu/>), a European consortium which aims to identify risk factors for NP.

Conflict of interest statement

B. H. Smith is a member of the DOLORisk consortium which is funded by the European Commission Horizon 2020 (ID: 633491) and is partly supported by this grant. He has received, on behalf of his institution, occasional lecture and consultancy fees from Pfizer Ltd, Napp Pharmaceuticals, Grunenthal and Eli Lilly. H. L. Hébert and A. Veluchamy are supported by DOLORisk. N. Torrance has no conflicts of interest to declare.

Article history:

Received 12 September 2016

Received in revised form 16 November 2016

Accepted 30 November 2016

Available online 7 December 2016

References

- Abbott CA, Malik RA, van Ross ER, Kulkarni J, Boulton AJ. Prevalence and characteristics of painful diabetic neuropathy in a large community-based diabetic population in the U.K. *Diabetes Care* 2011;34:2220–4.
- AlQuliti K. Predictors of painful diabetic neuropathy in Saudi patients with type 2 diabetes. *J Pain Relief* 2015;4:181.
- Barrett JC, Clayton DG, Concannon P, Akolkar B, Cooper JD, Erlich HA, Julier C, Morahan G, Nerup J, Nierras C, Plagnol V, Pociot F, Schuilenburg H, Smyth DJ, Stevens H, Todd JA, Walker NM, Rich SS. Genome-wide association study and meta-analysis find that over 40 loci affect risk of type 1 diabetes. *Nat Genet* 2009;41:703–7.
- Benbow SJ, Williams R, MacFarlane IA. Smoking habits and painful diabetic neuropathy. *J Diabetes Complications* 1997;11:334–7.
- Cheng KI, Lin SR, Chang LL, Wang JY, Lai CS. Association of the functional A118G polymorphism of OPRM1 in diabetic patients with foot ulcer pain. *J Diabetes Complications* 2010;24:102–8.
- Cortez J, Reis C, Cardoso Y, Onofre A, Piovezan AP. Prevalence of neuropathic pain and associated factors in diabetes mellitus type 2 patients seen in outpatient setting. *Revista Dor* 2014;15:256–9.
- D'Amato C, Morganti R, Greco C, Di Gennaro F, Cacciotti L, Longo S, Mataluni G, Lauro D, Marfia GA, Spallone V. Diabetic peripheral neuropathic pain is a stronger predictor of depression than other diabetic complications and comorbidities. *Diab Vasc Dis Res* 2016;13:418–28.
- Daousi C, MacFarlane IA, Woodward A, Nurmikko TJ, Bundred PE, Benbow SJ. Chronic painful peripheral neuropathy in an urban community: a controlled comparison of people with and without diabetes. *Diabet Med* 2004;21:976–82.
- Davies M, Brophy S, Williams R, Taylor A. The prevalence, severity, and impact of painful diabetic peripheral neuropathy in type 2 diabetes. *Diabetes Care* 2006;29:1518–22.
- DiBonaventura MD, Cappelleri JC, Joshi AV. Association between pain severity and health care resource use, health status, productivity and related costs in painful diabetic peripheral neuropathy patients. *Pain Med* 2011;12:799–807.
- Erbas T, Ertas M, Yucel A, Keskinaslan A, Senocak M, Group TS. Prevalence of peripheral neuropathy and painful peripheral neuropathy in Turkish diabetic patients. *J Clin Neurophysiol* 2011;28:51–5.
- Finnerup NB, Haroutounian S, Kamerman P, Baron R, Bennett DL, Bouhassira D, Cruccu G, Freeman R, Hansson P, Nurmikko T, Raja SN, Rice AS, Serra J, Smith BH, Treede RD, Jensen TS. Neuropathic pain: an updated grading system for research and clinical practice. *PAIN* 2016;157:1599–606.
- Forrest KY, Maser RE, Pambianco G, Becker DJ, Orchard TJ. Hypertension as a risk factor for diabetic neuropathy: a prospective study. *Diabetes* 1997;46:665–70.
- Galer BS, Gianas A, Jensen MP. Painful diabetic polyneuropathy: epidemiology, pain description, and quality of life. *Diabetes Res Clin Pract* 2000;47:123–8.
- Gore M, Brandenburg NA, Dukes E, Hoffman DL, Tai KS, Stacey B. Pain severity in diabetic peripheral neuropathy is associated with patient functioning, symptom levels of anxiety and depression, and sleep. *J Pain Symptom Manage* 2005;30:374–85.
- Haase CL, Tybjaerg-Hansen A, Nordestgaard BG, Frikke-Schmidt R. HDL cholesterol and risk of type 2 diabetes: a Mendelian Randomization Study. *Diabetes* 2015;64:3328–33.
- Halawa MR, Karawagh A, Zeidan A, Mahmoud AE, Sakr M, Hegazy A. Prevalence of painful diabetic peripheral neuropathy among patients suffering from diabetes mellitus in Saudi Arabia. *Curr Med Res Opin* 2010;26:337–43.
- Harris M, Eastman R, Cowie C. Symptoms of sensory neuropathy in adults with NIDDM in the U.S. population. *Diabetes Care* 1993;16:1446–52.
- Hyttinen V, Kaprio J, Kinnunen L, Koskenvuo M, Tuomilehto J. Genetic liability of type 1 diabetes and the onset age among 22,650 young Finnish twin pairs: a Nationwide Follow-Up Study. *Diabetes* 2003;52:1052–5.
- International Diabetes Federation. IDF Diabetes Atlas, 7th ed. Brussels: International Diabetes Federation, 2015. Available at: <http://www.diabetesatlas.org>. Accessed August 9, 2016.
- Jambart S, Ammache Z, Haddad F, Younes A, Hassoun A, Abdalla K, Selwan CA, Sunna N, Wajsbrot D, Yousef E. Prevalence of painful diabetic peripheral neuropathy among patients with diabetes mellitus in the Middle East region. *J Int Med Res* 2011;39:366–77.
- Jensen TS, Baron R, Haanpää M, Kalso E, Loeser JD, Rice AS, Treede RD. A new definition of neuropathic pain. *PAIN* 2011;152:2204–5.
- Li QS, Cheng P, Favis R, Wickenden A, Romano G, Wang H. SCN9A variants may be implicated in neuropathic pain associated with diabetic peripheral neuropathy and pain severity. *Clin J Pain* 2015;31:976–82.
- Li Y, Tong N. Angiotensin-converting enzyme I/D polymorphism and diabetic peripheral neuropathy in type 2 diabetes mellitus: a meta-analysis. *J Renin Angiotensin Aldosterone Syst* 2015;16:787–92.
- Meisinger C, Doring A, Thorand B, Heier M, Lowel H. Body fat distribution and risk of type 2 diabetes in the general population: are there differences between men and women? The MONICA/KORA Augsburg cohort study. *Am J Clin Nutr* 2006;84:483–9.
- Meng W, Deshmukh HA, Donnelly LA, Wellcome Trust Case Control Consortium 2 (WTCCC2), Surrogate markers for Micro- and Macro-vascular hard endpoints for Innovative diabetes Tools (SUMMIT) study group, Torrance N, Colhoun HM, Palmer CN, Smith BH. A genome-wide association study provides evidence of sex-specific involvement of chr1p35.1 (ZSCAN20-TLR12P) and chr8p23.1 (HMGBI1P46) with diabetic neuropathic pain. *EBioMedicine* 2015;2:1386–93.
- Meng W, Deshmukh HA, van Zuydam NR, Liu Y, Donnelly LA, Zhou K, Wellcome Trust Case Control Consortium 2, Surrogate Markers for Micro and Macro-Vascular Hard Endpoints for Innovative Diabetes Tools Study Group, Morris AD, Colhoun HM, Palmer CN, Smith BH. A genome-wide association study suggests an association of Chr8p21.3 (GFRA2) with diabetic neuropathic pain. *Eur J Pain* 2015;19:392–9.
- Miralles-Garcia JM, de Pablos-Velasco P, Cabrerizo L, Perez M, Lopez-Gomez V. Prevalence of distal diabetic polyneuropathy using quantitative sensory methods in a population with diabetes of more than 10 years' disease duration. *Endocrinol Nutr* 2010;57:414–20.
- Poulsen P, Kyvik KO, Vaag A, Beck-Nielsen H. Heritability of type II (non-insulin-dependent) diabetes mellitus and abnormal glucose tolerance—a population-based twin study. *Diabetologia* 1999;42:139–45.
- Purwata TE. High TNF-alpha plasma levels and macrophages iNOS and TNF-alpha expression as risk factors for painful diabetic neuropathy. *J Pain Res* 2011;4:169–75.
- Sladek R, Rocheleau G, Rung J, Dina C, Shen L, Serre D, Boutin P, Vincent D, Belisle A, Hadjadj S, Balkau B, Heude B, Charpentier G, Hudson TJ, Montpetit A, Pshezhetsky AV, Prentki M, Posner BI, Balding DJ,

- Meyre D, Polychronakos C, Froguel P. A genome-wide association study identifies novel risk loci for type 2 diabetes. *Nature* 2007;445:881–5.
- [32] Sorensen L, Molyneaux L, Yue DK. Insensate versus painful diabetic neuropathy: the effects of height, gender, ethnicity and glycaemic control. *Diabetes Res Clin Pract* 2002;57:45–51.
- [33] Spallone V, Morganti R, D'Amato C, Cacciotti L, Fedele T, Maiello MR, Marfia G. Clinical correlates of painful diabetic neuropathy and relationship of neuropathic pain with sensorimotor and autonomic nerve function. *Eur J Pain* 2011;15:153–60.
- [34] Tang TS, Prior SL, Li KW, Ireland HA, Bain SC, Hurel SJ, Cooper JA, Humphries SE, Stephens JW. Association between the rs1050450 glutathione peroxidase-1 (C > T) gene variant and peripheral neuropathy in two independent samples of subjects with diabetes mellitus. *Nutr Metab Cardiovasc Dis* 2012;22:417–25.
- [35] Tesfaye S. Recent advances in the management of diabetic distal symmetrical polyneuropathy. *J Diabetes Investig* 2011;2:33–42.
- [36] Themistocleous AC, Ramirez JD, Shillo PR, Lees JG, Selvarajah D, Orengo C, Tesfaye S, Rice ASC, Bennett DLH. The Pain in Neuropathy Study (PiNS): a cross-sectional observational study determining the somatosensory phenotype of painful and painless diabetic neuropathy. *PAIN* 2016;157:1132–45.
- [37] Tölle T, Xu X, Sadosky AB. Painful diabetic neuropathy: a cross-sectional survey of health state impairment and treatment patterns. *J Diabetes Complications* 2006;20:26–33.
- [38] Van Acker K, Bouhassira D, De Bacquer D, Weiss S, Matthys K, Raemen H, Mathieu C, Colin IM. Prevalence and impact on quality of life of peripheral neuropathy with or without neuropathic pain in type 1 and type 2 diabetic patients attending hospital outpatients clinics. *Diabetes Metab* 2009;35:206–13.
- [39] van Hecke O, Austin SK, Khan RA, Smith BH, Torrance N. Neuropathic pain in the general population: a systematic review of epidemiological studies. *PAIN* 2014;155:654–62.
- [40] van Hecke O, Kamerman PR, Attal N, Baron R, Bjornsdottir G, Bennett DL, Bennett MI, Bouhassira D, Diatchenko L, Freeman R, Freynhagen R, Haanpaa M, Jensen TS, Raja SN, Rice AS, Seltzer Z, Thorgeirsson TE, Yarnitsky D, Smith BH. Neuropathic pain phenotyping by international consensus (NeuroPPIC) for genetic studies: a NeuPSIG systematic review, Delphi survey, and expert panel recommendations. *PAIN* 2015;156:2337–53.
- [41] Wu EQ, Borton J, Said G, Le TK, Monz B, Rosilio M, Avoinet S. Estimated prevalence of peripheral neuropathy and associated pain in adults with diabetes in France. *Curr Med Res Opin* 2007;23:2035–42.
- [42] Ziegler D, Rathmann W, Dickhaus T, Meisinger C, Mielck A. Neuropathic pain in diabetes, prediabetes and normal glucose tolerance: the MONICA/KORA Augsburg Surveys S2 and S3. *Pain Med* 2009;10:393–400.
- [43] Ziegler D, Rathmann W, Meisinger C, Dickhaus T, Mielck A. Prevalence and risk factors of neuropathic pain in survivors of myocardial infarction with pre-diabetes and diabetes. The KORA Myocardial Infarction Registry. *Eur J Pain* 2009;13:582–7.